



A note on hypocoercivity for kinetic equations with heavy-tailed equilibrium

Nathalie Ayi, Maxime Herda, H  l  ne Hivert, Isabelle Tristani

► To cite this version:

Nathalie Ayi, Maxime Herda, H  l  ne Hivert, Isabelle Tristani. A note on hypocoercivity for kinetic equations with heavy-tailed equilibrium. *Comptes Rendus Math  matique*, Elsevier Masson, 2020, 358 (3), pp.333-340. 10.5802/crmath.46 . hal-02389146v2

HAL Id: hal-02389146

<https://hal.archives-ouvertes.fr/hal-02389146v2>

Submitted on 16 Mar 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin  e au d  p  t et    la diffusion de documents scientifiques de niveau recherche, publi  s ou non,   manant des   tablissements d'enseignement et de recherche fran  ais ou   trangers, des laboratoires publics ou priv  s.

A note on hypocoercivity for kinetic equations with heavy-tailed equilibrium

Nathalie Ayi ^a, Maxime Herda ^b, H       Hivert ^c and Isabelle Tristani ^d

^a*Sorbonne Universit  , Universit   de Paris, CNRS, Laboratoire Jacques-Louis Lions, 4 place Jussieu, 75005 Paris, France*

^b*Inria, Univ. Lille, CNRS, UMR 8524 - Laboratoire Paul Painlev  , F-59000 Lille, France*

^c*Univ. Lyon,   cole centrale de Lyon, CNRS UMR 5208, Institut Camille Jordan, F-69134   cully, France*

^d*DMA,   cole Normale Sup  rieure, CNRS, PSL Research University, 45 rue d'Ulm, 75005 Paris, France*

Received *****; accepted after revision ++++++

Presented by

Abstract

In this paper we are interested in the large time behavior of linear kinetic equations with heavy-tailed local equilibria. Our main contribution concerns the kinetic L  vy-Fokker-Planck equation, for which we adapt hypocoercivity techniques in order to show that solutions converge exponentially fast to the global equilibrium. Compared to the classical kinetic Fokker-Planck equation, the issues here concern the lack of symmetry of the non-local L  vy-Fokker-Planck operator and the understanding of its regularization properties. As a complementary related result, we also treat the case of the heavy-tailed BGK equation.

R  sum  

Une note sur l'hypocoercivit   pour les   quations cin  tiques avec   quilibres    queue lourde. Dans cet article, on s'int  resse au comportement en temps long d'  quations cin  tiques lin  aires dont les   quilibres locaux sont    queue lourde. Notre contribution principale concerne l'  quation de L  vy-Fokker-Planck cin  tique, pour laquelle nous adaptons des techniques d'hypocoercivit   afin de d  montrer la convergence exponentielle des solutions vers un   quilibre global. En comparant au cas de l'  quation de Fokker-Planck cin  tique classique, les enjeux ici sont li  s au manque de sym  trie de l'op  rateur non-local de L  vy-Fokker-Planck et    la compr  hension de ses propri  t  s de r  gularisation. En compl  ment de notre analyse, nous traitons   galement le cas de l'  quation de BGK    queue lourde.

Email addresses: nathalie.ayi@sorbonne-universite.fr (Nathalie Ayi), maxime.herda@inria.fr (Maxime Herda), helene.hivert@ec-lyon.fr (H       Hivert), isabelle.tristani@ens.fr (Isabelle Tristani).

1. Introduction

We consider a distribution function $f \equiv f(t, x, v)$ depending on time $t \geq 0$, position $x \in \mathbb{T}^d$ and velocity $v \in \mathbb{R}^d$ which satisfies the fractional kinetic Fokker-Planck equation

$$\partial_t f + v \cdot \nabla_x f = \nabla_v \cdot (vf) - (-\Delta_v)^{\alpha/2} f. \quad (1)$$

Here we assume $\alpha \in (0, 2)$ and the fractional Laplacian $-(-\Delta_v)^{\alpha/2}$ is such that for any Schwartz function $g : \mathbb{R}^d \rightarrow \mathbb{R}$, one has $\mathcal{F}((- \Delta_v)^{\alpha/2} g)(\xi) = |\xi|^\alpha \mathcal{F}(g)(\xi)$ where $\mathcal{F}(\cdot)$ denotes the Fourier transform. There are many equivalent definitions of the fractional Laplacian (see [12]). Among them we shall use

$$(-\Delta_v)^{\alpha/2} g(v) = C_{d,\alpha} \text{P.V.} \int_{\mathbb{R}^d} \frac{g(v) - g(w)}{|v - w|^{d+\alpha}} dw, \quad (2)$$

where P.V. stands for the principal value and the constant is given by $C_{d,\alpha} = 2^\alpha \Gamma(\frac{d+\alpha}{2}) / (\pi^{d/2} |\Gamma(-\frac{\alpha}{2})|)$ where $\Gamma(\cdot)$ is the Gamma function. In the following we drop the principal value in the notations. We denote the Lévy-Fokker-Planck operator appearing on the right-hand side of (1) by

$$L_\alpha g = \nabla_v \cdot (vg) - (-\Delta_v)^{\alpha/2} g.$$

By passing to Fourier variables one has $\mathcal{F}(L_\alpha g)(\xi) = -\xi \cdot \nabla_\xi \hat{g}(\xi) - |\xi|^\alpha \hat{g}(\xi)$, where $\hat{g} = \mathcal{F}(g)$. From this formula, one sees that the function

$$\mu_\alpha(v) = Z_{d,\alpha}^{-1} \mathcal{F}^{-1} \left(e^{-|\xi|^\alpha / \alpha} \right)$$

with $Z_{d,\alpha}$ chosen such that $\int \mu_\alpha = 1$ is a probability distribution such that $L_\alpha \mu_\alpha = 0$. Observe that away from the origin, the Fourier transform of μ_α is smooth and rapidly decaying at infinity. The singularity at $\xi = 0$ behaves like $|\xi|^\alpha$ at principal order which yields that $\mu_\alpha(v)$ should decay as $|v|^{-\alpha-d}$ when $|v| \rightarrow \infty$. Actually, one has the following more precise bounds coming from [4, Theorem 3.1] and references in the proof (see also the references in [1]). There are positive constants $C_1 = C_1(\alpha, d) > 0$ and $C_2 = C_2(\alpha, d) > 0$ such that for all $v \in \mathbb{R}^d$ one has

$$C_1^{-1} \leq (|v|^{d+\alpha} + 1) \mu_\alpha(v) \leq C_1, \quad (3)$$

and

$$C_2^{-1} |v| \leq (|v|^{2+d+\alpha} + 1) |\nabla_v \mu_\alpha(v)| \leq C_2 |v|. \quad (4)$$

In the following, given some measurable non-negative function $\nu \equiv \nu(v)$ we denote by $L_v^2(\nu)$ and $L_{x,v}^2(\nu)$ the spaces of measurable functions g of respectively the v and the (x, v) variables such that $|g|^2 \nu$ is integrable. We endow these spaces with their canonical scalar product and norm. We also introduce the corresponding Sobolev space $H_{x,v}^1(\nu)$ associated with the norm

$$\|g\|_{H_{x,v}^1(\nu)}^2 = \|g\|_{L_{x,v}^2(\nu)}^2 + \|\nabla_x g\|_{L_{x,v}^2(\nu)}^2 + \|\nabla_v g\|_{L_{x,v}^2(\nu)}^2.$$

Finally given an integrable function g , we denote $\langle g \rangle = \iint_{\mathbb{T}^d \times \mathbb{R}^d} g(x, v) dv dx$ the global mass of g . The main result of this paper is the following.

Theorem 1.1 *Let f solve the kinetic Lévy-Fokker-Planck equation (1) with initial data $f^{in} \in H_{x,v}^1(\mu_\alpha^{-1})$. Then, for all $t \geq 0$ one has*

$$\|f(t) - \langle f^{in} \rangle \mu_\alpha\|_{H_{x,v}^1(\mu_\alpha^{-1})} \leq C \|f^{in} - \langle f^{in} \rangle \mu_\alpha\|_{H_{x,v}^1(\mu_\alpha^{-1})} e^{-\lambda t}$$

for some constant $C \geq 1$ and $\lambda > 0$ depending only on d and α .

Let us mention that these results have been obtained as a preliminary step towards the conception and analysis of numerical schemes preserving the long-time behavior of these equations. This topic is an ongoing work [2] in the spirit of what has previously been done in [8,3] in the case of the classical Fokker-Planck equation. The compatibility of our schemes with anomalous diffusion limit will also be investigated (see [7] for more details).

Before going into the analysis of our problem, let us recall that results on large time behavior of solutions to the homogeneous version of (1), namely $\partial_t f(t, v) = L_\alpha f(t, v)$, have been obtained in [9] in spaces of type $L_v^2(\mu_\alpha^{-1})$ (among others) and later in [13] in larger Lebesgue spaces. Notice that the presence of the transport operator in our equation (1) makes the analysis more intricate and requires the use of hypocoercivity techniques. In the present note, we use H^1 type hypocoercivity as presented in [14] or [10] for example. Note also that fractional hypocoercivity has already been studied recently in [5] where a L^2 -hypocoercivity approach is developed. In this sense, their framework is quite different, note however that it is also more general than ours (in terms of phase space and linear operators). Their results in particular imply an exponential convergence towards equilibrium in the torus in L^2 for our models.

In the same spirit of our work, let us also mention the paper [11] in which some hypoelliptic estimates are obtained on the non homogeneous fractional Kolmogorov equation (there is no drift term in the studied equation). The method of proof is quite close (based on the use of weighted Lyapunov functional) but the final goal is different in the latter since the main concern is about regularization properties of the equation and not convergence towards the equilibrium.

In the present study we focus on a good understanding of the structure of the Lévy-Fokker-Planck operator since we endeavour to carry out our computations as simply as possible in order to adapt our analysis to a discrete framework in [2]. In particular let us point out that we do not need fractional derivatives in our Lyapunov functionals and our proof does not rely on Fourier transform. In this sense our method differs completely from that of [11] and the recent [5] in which a mode by mode analysis is developed.

Outline of the note. From Section 2 to Section 4, we carry out the analysis of the properties of the Lévy-Fokker-Planck operator that will be useful for proving our main result. Then, the proof of Theorem 1.1 is done in Section 5. In the last section we state and prove the equivalent of Theorem 1.1 for the BGK equation with heavy-tailed equilibrium.

Notations. For simplicity, in the subsequent proofs, we denote by C a positive constant depending only on fixed numbers (including d and α) and its value may change from line to line.

2. The Lévy-Fokker-Planck operator as bilinear form

The following quite simple decomposition is actually one of the key elements of our hypocoercive analysis carried out in Section 5. Compared to the non-fractional case, we here have a lack of symmetry of our operator in $L_v^2(\mu_\alpha^{-1})$ and the following splitting is very helpful to simplify the computations. Moreover, in the non-fractional case, there is a gain of weight in velocity which comes from the particular form of the gradient of the Gaussian equilibrium. Even though we no longer have such a gain in our case, we are still able to close our estimates thanks to the following splitting.

Proposition 2.1 *One has the decomposition*

$$-\langle L_\alpha f, g \rangle_{L_v^2(\mu_\alpha^{-1})} = \mathcal{S}_v(f, g) + \mathcal{A}_v(f, g),$$

where \mathcal{S}_v and \mathcal{A}_v are bilinear forms that are respectively symmetric and skew-symmetric and defined by

$$\mathcal{S}_v(f, g) = \frac{C_{d,\alpha}}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^d} \frac{[(f\mu_\alpha^{-1})(v) - (f\mu_\alpha^{-1})(w)][(g\mu_\alpha^{-1})(v) - (g\mu_\alpha^{-1})(w)]}{|v - w|^{d+\alpha}} \mu_\alpha(v) \, dw \, dv,$$

and

$$\begin{aligned} \mathcal{A}_v(f, g) &= \frac{C_{d,\alpha}}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^d} \frac{(f\mu_\alpha^{-1})(w)(g\mu_\alpha^{-1})(v) - (f\mu_\alpha^{-1})(v)(g\mu_\alpha^{-1})(w)}{|v - w|^{d+\alpha}} \mu_\alpha(v) \, dw \, dv \\ &\quad + \frac{1}{2} \int_{\mathbb{R}^d} (f v \cdot \nabla_v (g\mu_\alpha^{-1}) - g v \cdot \nabla_v (f\mu_\alpha^{-1})) \, dv, \end{aligned}$$

where $C_{d,\alpha}$ is defined in (2).

We skip the proof of this proposition since it is based on simple computations using the formula (2), integration by parts and the fact that $L_\alpha \mu_\alpha = 0$.

Observe that a direct consequence of the Cauchy-Schwarz inequality is

$$\mathcal{S}_v(f, g) \leq \mathcal{S}_v(f, f)^{1/2} \mathcal{S}_v(g, g)^{1/2}, \quad (5)$$

for $f, g \in D(L_\alpha)$. Moreover, the symmetric form \mathcal{S}_v is non-negative and $\mathcal{S}_v(f, f)$ vanishes when $f\mu_\alpha^{-1}$ is constant. This yields that the nullspace of L_α is exactly given by $\mathbb{R}\mu_\alpha$. From there the orthogonal projection Π onto the nullspace of L_α is given by

$$(\Pi g)(v) = \left(\int_{\mathbb{R}^d} g(w) \, dw \right) \mu_\alpha(v).$$

3. Coercivity results for the Lévy-Fokker-Planck operator

One has the following coercivity result taken from [9, Theorem 2] and originating from [6]. While the previous references derive the inequality via a semigroup approach, let us mention that an elementary analytical proof is given by Wang [15] and came to our attention thanks to [5].

Lemma 3.1 ([6], [9], [15]...) *There is a constant $C_P \equiv C_P(\alpha, d) > 0$ such that for all $f \in D(L_\alpha)$,*

$$\|f - \Pi f\|_{L_v^2(\mu_\alpha^{-1})}^2 \leq C_P \mathcal{S}_v(f, f). \quad (6)$$

We now show that the dissipation $\mathcal{S}_v(f, f)$ also provides some fractional Sobolev regularity. We introduce the fractional Sobolev space H_v^s with $s \in (0, 1)$ with norm defined by $\|\cdot\|_{H_v^s}^2 = \|\cdot\|_{L_v^2}^2 + \|\cdot\|_{\dot{H}_v^s}^2$ where the homogeneous Sobolev norm is given by $\|g\|_{\dot{H}_v^s}^2 := \|(-\Delta)^{s/2} g\|_{L_v^2}^2$. One can prove that there exists a positive constant $\tilde{C}_{d,s}$ such that

$$\|g\|_{\dot{H}_v^s}^2 = \tilde{C}_{d,s} \iint_{\mathbb{R}^d \times \mathbb{R}^d} |f(v) - f(w)|^2 |v - w|^{-(d+2s)} \, dw \, dv. \quad (7)$$

Lemma 3.2 *There exists $C_R \equiv C_R(\alpha, d) > 0$ such that for all $f \in D(L_\alpha)$,*

$$\mathcal{S}_v(f, f) \geq C_R^{-1} \left(\|f\mu_\alpha^{-1/2}\|_{\dot{H}_v^{\alpha/2}}^2 - \|f\mu_\alpha^{-1/2}\|_{L_v^2}^2 \right).$$

Proof. Using that $(a + b)^2 \geq a^2/2 - b^2$, we have

$$\mathcal{S}_v(f, f) \geq \frac{C_{d,\alpha}}{2} \iint_{|v-w| \leq 1} \frac{|(\mu_\alpha^{-1}f)(v) - (\mu_\alpha^{-1}f)(w)|^2}{|v - w|^{d+\alpha}} \mu_\alpha(v) \, dw \, dv \geq \frac{C_{d,\alpha}}{2} \left(\frac{1}{2} I_1 - I_2 \right).$$

The first term is

$$\begin{aligned}
I_1 &= \iint_{|v-w| \leq 1} \frac{|(\mu_\alpha^{-1/2}f)(v) - (\mu_\alpha^{-1/2}f)(w)|^2}{|v-w|^{d+\alpha}} dw dv \\
&= \tilde{C}_{d, \frac{\alpha}{2}}^{-1} \|\mu_\alpha^{-1/2}f\|_{\dot{H}_v^{\alpha/2}}^2 - \iint_{|v-w| \geq 1} \frac{|(\mu_\alpha^{-1/2}f)(v) - (\mu_\alpha^{-1/2}f)(w)|^2}{|v-w|^{d+\alpha}} dw dv \\
&\geq \tilde{C}_{d, \frac{\alpha}{2}}^{-1} \|\mu_\alpha^{-1/2}f\|_{\dot{H}_v^{\alpha/2}}^2 - C \|\mu_\alpha^{-1/2}f\|_{L_v^2}^2
\end{aligned}$$

where $\tilde{C}_{d, \frac{\alpha}{2}}$ is defined in (7) and for the last inequality, we used the integrability of $|v-w|^{-d-\alpha} \mathbf{1}_{|v-w| \geq 1}$, once in v and once in w . The second term is

$$I_2 = \iint_{|v-w| \leq 1} \frac{|\mu_\alpha^{1/2}(v) - \mu_\alpha^{1/2}(w)|^2}{|v-w|^{d+\alpha}} |f(w)|^2 |\mu_\alpha^{-1}(w)|^2 dw dv.$$

To treat I_2 , we use Taylor formula to write

$$I_2 = \iint_{|w| \leq 1} \frac{\left| \int_0^1 \nabla(\mu_\alpha^{1/2})(v - \theta w) \cdot w d\theta \right|^2}{|w|^{d+\alpha}} \mu_\alpha^{-1}(v-w) |f(v-w) \mu_\alpha^{-1/2}(v-w)|^2 dw dv.$$

Performing now the changes of variables $v \rightarrow v - \theta w$ and then $\theta \rightarrow 1 - \theta$, we get:

$$I_2 \leq \iint_{|w| \leq 1} \int_0^1 \frac{|\nabla(\mu_\alpha^{1/2})(v)|^2}{|w|^{d+\alpha-2}} \mu_\alpha^{-1}(v - \theta w) |f(v - \theta w) \mu_\alpha^{-1/2}(v - \theta w)|^2 d\theta dw dv.$$

Notice that, using (3) and since $|w| \leq 1$, we have $\mu_\alpha^{-1}(v - \theta w) \leq C(1 + |v|^{d+\alpha})$. Then, using (4), one can prove that $|\nabla(\mu_\alpha^{1/2})(v)|^2 \mu_\alpha^{-1}(v - \theta w) \leq C$. Consequently, we obtain

$$I_2 \leq C \iint_{|w| \leq 1} \int_0^1 \frac{1}{|w|^{d+\alpha-2}} |f(v - \theta w) \mu_\alpha^{-1/2}(v - \theta w)|^2 d\theta dw dv$$

and thus performing a change of variable $I_2 \leq C \|f \mu_\alpha^{-1/2}\|_{L_v^2}^2$. This ends the proof. \square

Proposition 3.3 *There is $C_F \equiv C_F(\alpha, d)$ such that for all $f \in D(L_\alpha)$,*

$$\|(f - \Pi f) \mu_\alpha^{-1/2}\|_{\dot{H}_v^{\alpha/2}}^2 \leq C_F \mathcal{S}_v(f, f). \quad (8)$$

Proof. Let us now summarize the estimates that we have obtained in the two previous lemma. We have $\mathcal{S}_v(f, f) \geq C_P^{-1} \|f - \Pi f\|_{L_v^2(\mu_\alpha^{-1/2})}^2$ and $\mathcal{S}_v(f, f) \geq C_R^{-1} (\|f \mu_\alpha^{-1/2}\|_{\dot{H}_v^{\alpha/2}}^2 - \|f \mu_\alpha^{-1/2}\|_{L_v^2}^2)$. Moreover, one can notice that $\mathcal{S}_v(f, f) = \mathcal{S}_v(f - \Pi f, f - \Pi f)$. As a consequence, an appropriate convex combination of the two previous inequalities shows (8). \square

4. An interpolation inequality

In this section we prove an interpolation result which is crucial in the proof of Theorem 1.1.

Proposition 4.1 *For all $\varepsilon > 0$, there is $K(\varepsilon) \equiv K(\varepsilon, \alpha, d) > 0$ such that*

$$\|\nabla_v f\|_{L_v^2(\mu_\alpha^{-1})}^2 \leq K(\varepsilon) \left(\mathcal{S}_v(f, f) + \|\Pi f\|_{L_v^2(\mu_\alpha^{-1})}^2 \right) + \varepsilon C_F \mathcal{S}_v(\nabla_v f, \nabla_v f) \quad (9)$$

where the constant C_F is defined in Proposition 3.3.

Proof. One can use the chain rule and an interpolation of \dot{H}_v^1 between $\dot{H}_v^{\alpha/2}$ and $\dot{H}_v^{1+\alpha/2}$ (easily shown in Fourier variables) to get

$$\begin{aligned}
\|(\nabla_v f)\mu_\alpha^{-1/2}\|_{L_v^2}^2 &\leq 2\|\nabla_v(f\mu_\alpha^{-1/2})\|_{L_v^2}^2 + 2\|f(\nabla_v\mu_\alpha^{-1/2})\|_{L_v^2}^2 \\
&\leq K(\varepsilon)\|f\mu_\alpha^{-1/2}\|_{\dot{H}_v^{\alpha/2}}^2 + \varepsilon\|\nabla_v(f\mu_\alpha^{-1/2})\|_{\dot{H}_v^{\alpha/2}}^2 + 2\|f(\nabla_v\mu_\alpha^{-1/2})\|_{L_v^2}^2 \\
&\leq K(\varepsilon)\|f\mu_\alpha^{-1/2}\|_{H_v^{\alpha/2}}^2 + \varepsilon\|(\nabla_v f)\mu_\alpha^{-1/2}\|_{H_v^{\alpha/2}}^2 + C\|f\mu_\alpha^{-1/2}\|_{L_v^2}^2 \\
&\leq K(\varepsilon)\|f\mu_\alpha^{-1/2}\|_{H_v^{\alpha/2}}^2 + \varepsilon\|(\nabla_v f)\mu_\alpha^{-1/2}\|_{H_v^{\alpha/2}}^2
\end{aligned}$$

up to changing the value of $K(\varepsilon)$ and where we used the fact that $|(\nabla_v\mu_\alpha)\mu_\alpha^{-1}| \in L^\infty(\mathbb{R}^d)$ to bound the third term. Now observe that

$$\|f\mu_\alpha^{-1/2}\|_{H_v^{\alpha/2}}^2 \leq 2\left(\|(f - \Pi f)\mu_\alpha^{-1/2}\|_{H_v^{\alpha/2}}^2 + \|(\Pi f)\mu_\alpha^{-1/2}\|_{H_v^{\alpha/2}}^2\right),$$

and that $\|(\Pi f)\mu_\alpha^{-1/2}\|_{H_v^{\alpha/2}} = \|\Pi f\|_{L_v^2(\mu_\alpha^{-1})}\|\mu_\alpha^{1/2}\|_{H_v^{\alpha/2}}$ with $\|\mu_\alpha^{1/2}\|_{H_v^{\alpha/2}} \leq C$ since $\mu_\alpha^{1/2} \in H_v^1$ from (3) and (4). Moreover, one has $\nabla_v f = \nabla_v \tilde{f} - \Pi \nabla_v f$. One can conclude by using (8) twice. \square

5. Proof of Theorem 1.1

Up to changing f^{in} by $f^{\text{in}} - \langle f^{\text{in}} \rangle \mu_\alpha$, we assume that $\iint_{\mathbb{T}^d \times \mathbb{R}^d} f(t, x, v) dv dx = 0$ at $t = 0$, so that by conservation it also holds for all time $t > 0$. We introduce a new norm on the weighted Sobolev space $H_{x,v}^1(\mu_\alpha^{-1})$. It is defined by

$$\|f\|^2 = \|f\|_{L_{x,v}^2(\mu_\alpha^{-1})}^2 + a\|\nabla_x f\|_{L_{x,v}^2(\mu_\alpha^{-1})}^2 + b\|\nabla_v f\|_{L_{x,v}^2(\mu_\alpha^{-1})}^2 + 2c\langle \nabla_x f, \nabla_v f \rangle_{L_{x,v}^2(\mu_\alpha^{-1})}, \quad (10)$$

where a, b and c are positive constants to be determined later on. Observe that as soon as $c^2 < ab$, one has that $\|\cdot\|$ is equivalent to $\|\cdot\|_{H_{x,v}^1(\mu_\alpha^{-1})}$. Let us note that the commutators $[\nabla_x, v \cdot \nabla_x]$ and $[\nabla_x, L_\alpha]$ vanish while $[\nabla_v, v \cdot \nabla_x] = \nabla_x$ and $[\nabla_v, L_\alpha] = \nabla_v$. Also observe that $v \cdot \nabla_x$ is skew-symmetric in $L_{x,v}^2(\mu_\alpha^{-1})$. Let us estimate the evolution of each term appearing in the new norm defined in (10) for f a solution of (1) with initial data f^{in} satisfying $\langle f^{\text{in}} \rangle = 0$. In the following the notation $\mathcal{S}_{x,v}$ denotes the integral of \mathcal{S}_v in the x variable. One has

$$\begin{aligned}
\frac{1}{2} \frac{d}{dt} \|f\|_{L_{x,v}^2(\mu_\alpha^{-1})}^2 &= -\mathcal{S}_{x,v}(f, f), \\
\frac{1}{2} \frac{d}{dt} \|\nabla_x f\|_{L_{x,v}^2(\mu_\alpha^{-1})}^2 &= -\mathcal{S}_{x,v}(\nabla_x f, \nabla_x f), \\
\frac{1}{2} \frac{d}{dt} \|\nabla_v f\|_{L_{x,v}^2(\mu_\alpha^{-1})}^2 &= -\mathcal{S}_{x,v}(\nabla_v f, \nabla_v f) + \|\nabla_v f\|_{L_{x,v}^2(\mu_\alpha^{-1})}^2 - \langle \nabla_x f, \nabla_v f \rangle_{L_{x,v}^2(\mu_\alpha^{-1})}, \\
\frac{d}{dt} \langle \nabla_x f, \nabla_v f \rangle_{L_{x,v}^2(\mu_\alpha^{-1})} &= -\|\nabla_x f\|_{L_{x,v}^2(\mu_\alpha^{-1})}^2 - 2\mathcal{S}_{x,v}(\nabla_x f, \nabla_v f) + \langle \nabla_x f, \nabla_v f \rangle_{L_{x,v}^2(\mu_\alpha^{-1})}.
\end{aligned}$$

Notice here that the keystone of the proof of the last equality is the splitting obtained in Proposition 2.1 and the Hilbertian setting. Indeed given any $g = e^{tL_\alpha} g_0$ and operators A and B , one has formally that $\frac{d}{dt} \langle Ag, Bg \rangle = \langle [A, L_\alpha]g, Bg \rangle + \langle Ag, [B, L_\alpha]g \rangle - 2\mathcal{S}_{x,v}(Ag, Bg)$. Therefore the skew symmetric part of the operator L_α only appears in commutators. This observation enables us to avoid loss of moments

in velocities in forthcoming estimates which one would face with bad rearrangements of the terms. By gathering all the previous estimates one gets

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|f\|^2 &= -\mathcal{S}_{x,v}(f, f) - a\mathcal{S}_{x,v}(\nabla_x f, \nabla_x f) - b\mathcal{S}_{x,v}(\nabla_v f, \nabla_v f) - c\|\nabla_x f\|_{L^2_{x,v}(\mu_\alpha^{-1})}^2 \\ &\quad + b\|\nabla_v f\|_{L^2_{x,v}(\mu_\alpha^{-1})}^2 - b\langle \nabla_x f, \nabla_v f \rangle_{L^2_{x,v}(\mu_\alpha^{-1})} \\ &\quad - 2c\mathcal{S}_{x,v}(\nabla_x f, \nabla_v f) + c\langle \nabla_x f, \nabla_v f \rangle_{L^2_{x,v}(\mu_\alpha^{-1})}. \end{aligned}$$

The first four terms are dissipation terms and the last four terms are remainder terms. Let us control the latter by the former ones. By integrating (5) in x and using Young's inequality one gets

$$|2c\mathcal{S}_{x,v}(\nabla_x f, \nabla_v f)| \leq \frac{2c^2}{b}\mathcal{S}_{x,v}(\nabla_x f, \nabla_x f) + \frac{b}{2}\mathcal{S}_{x,v}(\nabla_v f, \nabla_v f).$$

Then since $\int \nabla_v f dv = 0$, one has

$$\begin{aligned} b\left|\langle \nabla_x f, \nabla_v f \rangle_{L^2_{x,v}(\mu_\alpha^{-1})}\right| &= b\left|\langle \nabla_x f - \Pi(\nabla_x f), \nabla_v f \rangle_{L^2_{x,v}(\mu_\alpha^{-1})}\right| \\ &\leq bC_P^{1/2}\mathcal{S}_{x,v}(\nabla_x f, \nabla_x f)^{1/2}\|\nabla_v f\|_{L^2_{x,v}(\mu_\alpha^{-1})} \\ &\leq \frac{bC_P}{2}\mathcal{S}_{x,v}(\nabla_x f, \nabla_x f) + \frac{b}{2}\|\nabla_v f\|_{L^2_{x,v}(\mu_\alpha^{-1})}^2, \end{aligned}$$

where we used (6). Similarly

$$c\left|\langle \nabla_x f, \nabla_v f \rangle_{L^2_{x,v}(\mu_\alpha^{-1})}\right| \leq \frac{c^2C_P}{2b}\mathcal{S}_{x,v}(\nabla_x f, \nabla_x f) + \frac{b}{2}\|\nabla_v f\|_{L^2_{x,v}(\mu_\alpha^{-1})}^2.$$

For the last remainder term we use (9) integrated in x , namely

$$\|\nabla_v f\|_{L^2_{x,v}(\mu_\alpha^{-1})}^2 \leq K(\varepsilon)\left(\mathcal{S}_{x,v}(f, f) + \|\Pi f\|_{L^2_{x,v}(\mu_\alpha^{-1})}^2\right) + \varepsilon C_F\mathcal{S}_{x,v}(\nabla_v f, \nabla_v f).$$

We can use the Poincaré inequality on the torus (since f is mean-free) and the Jensen inequality to get $\|\Pi f\|_{L^2_{x,v}(\mu_\alpha^{-1})}^2 \leq \tilde{C}_P\|\nabla_x f\|_{L^2_{x,v}(\mu_\alpha^{-1})}^2$ where $\tilde{C}_P \equiv \tilde{C}_P(d)$ is the Poincaré constant of the d -dimensional torus. Thus eventually, one has

$$\frac{1}{2} \frac{d}{dt} \|f\|^2 + D(f, f) \leq 0, \quad (11)$$

where the dissipation is given by

$$\begin{aligned} D(f, f) &= (1 - 2bK(\varepsilon))\mathcal{S}_{x,v}(f, f) + \left(a - \frac{c^2}{b}\left(2 + \frac{C_P}{2}\right) - \frac{bC_P}{2}\right)\mathcal{S}_{x,v}(\nabla_x f, \nabla_x f) \\ &\quad + \left(\frac{b}{2} - 2b\varepsilon C_F\right)\mathcal{S}_{x,v}(\nabla_v f, \nabla_v f) + \left(c - 2b\tilde{C}_PK(\varepsilon)\right)\|\nabla_x f\|_{L^2_{x,v}(\mu_\alpha^{-1})}^2. \end{aligned}$$

Now choose consecutively ε, b, c and a such that $0 < \varepsilon < 1/(4C_F)$, $0 < b < 1/(2K(\varepsilon))$, $c > 2b\tilde{C}_PK(\varepsilon)$ and finally a large enough so that $a > c^2(2 + C_P/2)/b + bC_P/2$. It yields that the dissipation is non-negative and even that there is a constant $\lambda > 0$ (depending on a, b, c, ε) such that $D(f, f) \geq \lambda\|f\|^2$. By a Gronwall type argument we have that $\|f(t)\|$ decays exponentially to 0 when $t \rightarrow \infty$. \square

6. The case of the heavy-tailed BGK equation

In this last section we consider another simple kinetic model

$$\partial_t f + v \cdot \nabla_x f = \Pi_M f - f, \quad \text{with } (\Pi_M f)(t, x, v) = M(v) \int_{\mathbb{R}^d} f(t, x, w) dw, \quad (12)$$

for which the local equilibrium satisfies the following assumptions

$$M(v) > 0, \quad \int_{\mathbb{R}^d} M = 1, \quad \text{and } \nabla_v \ln(M) \in L^\infty. \quad (13)$$

This allows for heavy-tailed distributions, namely M such that $M(v) \sim_{|v| \rightarrow \infty} |v|^{-d-\alpha}$ with $\alpha \in (0, 2)$.

Theorem 6.1 *Assume that (13) holds and let f solve the BGK equation (12) starting from the initial data $f^{in} \in H_{x,v}^1(M^{-1})$. Then, for all $t \geq 0$ one has*

$$\|f(t) - \langle f^{in} \rangle M\|_{H_{x,v}^1(M^{-1})} \leq C \|f^{in} - \langle f^{in} \rangle M\|_{H_{x,v}^1(M^{-1})} e^{-\lambda t}$$

for some constant $C \geq 1$ and $\lambda > 0$ depending only on d and $\|\nabla_v \ln(M)\|_{L^\infty}$.

The proof is similar and simpler than that of Theorem 1.1. We skip many details as the reader may go back to the proof of Theorem 1.1 in order to recover them.

Proof of Theorem 6.1. Consider f a solution to (12) with initial data f^{in} satisfying $\langle f^{in} \rangle = 0$. Let us observe that the commutators $[\nabla_x, v \cdot \nabla_x]$ and $[\nabla_x, \Pi_M]$ vanish while $[\nabla_v, v \cdot \nabla_x] = \nabla_x$ and also $[\nabla_v, \Pi_M] = \nabla_v \ln(M) \Pi_M$. Now with this in mind, and defining the triple norm of f as in (10) with μ_α replaced by M , one gets

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|f\|^2 &= -\|f - \Pi_M f\|_{L_{x,v}^2(M^{-1})}^2 - a \|\nabla_x f - \Pi_M \nabla_x f\|_{L_{x,v}^2(M^{-1})}^2 - b \|\nabla_v f\|_{L_{x,v}^2(M^{-1})}^2 - c \|\nabla_x f\|_{L_{x,v}^2(M^{-1})}^2 \\ &\quad + b \langle \nabla_v \ln(M) \Pi_M f, \nabla_v f \rangle_{L_{x,v}^2(M^{-1})} - b \langle \nabla_x f, \nabla_v f \rangle_{L_{x,v}^2(M^{-1})} \\ &\quad - 2c \langle \nabla_x f, \nabla_v f \rangle_{L_{x,v}^2(M^{-1})} + c \langle \nabla_v \ln(M) \Pi_M f, \nabla_x f \rangle_{L_{x,v}^2(M^{-1})}. \end{aligned}$$

First, we notice that $\langle \nabla_x f, \nabla_v f \rangle_{L_{x,v}^2(M^{-1})} = \langle \nabla_x f - \Pi_M \nabla_x f, \nabla_v f \rangle_{L_{x,v}^2(M^{-1})}$ to deal with the third and fourth remainder terms with Cauchy-Schwarz inequality. The last remainder term requires some special care. Indeed, observe that since $\langle \nabla_v \ln(M) \Pi_M f, \Pi_M g \rangle$ vanishes for any g , one thus has

$$\langle \nabla_v \ln(M) \Pi_M f, \nabla_x f \rangle_{L_{x,v}^2(M^{-1})} \leq \|\nabla_v \ln(M)\|_{L^\infty} \|\Pi_M f\|_{L_{x,v}^2(M^{-1})} \|\nabla_x f - \Pi_M \nabla_x f\|_{L_{x,v}^2(M^{-1})}.$$

We also have that

$$\langle \nabla_v \ln(M) \Pi_M f, \nabla_v f \rangle_{L_{x,v}^2(M^{-1})} \leq \|\nabla_v \ln(M)\|_{L^\infty} \|\Pi_M f\|_{L_{x,v}^2(M^{-1})} \|\nabla_v f\|_{L_{x,v}^2(M^{-1})}.$$

Finally, we recall that $\|\Pi_M f\|_{L_{x,v}^2(M^{-1})} \leq \tilde{C}_P \|\nabla_x f\|_{L_{x,v}^2(M^{-1})}$ with \tilde{C}_P the Poincaré constant of the d -dimensional torus. Then using four times Young's inequality with well chosen weights, one obtains (11) with the dissipation

$$\begin{aligned} D(f, f) &= \|f - \Pi_M f\|_{L_{x,v}^2(M^{-1})}^2 + (a - b - 4c^2/b - C_M c/2) \|\nabla_x f - \Pi_M \nabla_x f\|_{L_{x,v}^2(M^{-1})}^2 \\ &\quad + (b - b/4 - b/4 - b/4) \|\nabla_v f\|_{L_{x,v}^2(M^{-1})}^2 + (c - bC_M - c/2) \|\nabla_x f\|_{L_{x,v}^2(M^{-1})}^2 \end{aligned}$$

with $C_M = \|\nabla_v \ln(M)\|_{L^\infty}^2 \tilde{C}_P^2$. One concludes as in Theorem 1.1 after choosing any $b > 0$, $c > 2bC_M$ and finally $a > 4c^2/b + b + C_{d,M}c/2$. \square

Acknowledgements

Maxime Herda thanks the LabEx CEMPI (ANR-11-LABX-0007-01). H       Hivert thanks the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (ERC starting grant MESOPROBIO n   639638). Isabelle Tristani thanks the ANR EFI: ANR-17-CE40-0030 and the ANR SALVE: ANR-19-CE40-0004 for their support.

This work is part of a collaborative research project that was initiated for the Junior Trimester Program in Kinetic Theory at the Hausdorff Research Institute for Mathematics in Bonn. Part of the work was carried out during the time spent at the institute. The authors are grateful for this opportunity and warmly acknowledge the HIM for the financial support and the hospitality they benefited during their stay.

References

- [1] Pedro Aceves-Sanchez and Ludovic Cesbron. Fractional Diffusion Limit for a Fractional Vlasov–Fokker–Planck Equation. *SIAM J. Math. Anal.* 51(1):469–488, 2019.
- [2] Nathalie Ayi, Maxime Herda, H       Hivert and Isabelle Tristani. On discrete fractional Fokker-Planck equations. *In preparation*.
- [3] Marianne Bessemoulin-Chatard, Maxime Herda, and Thomas Rey. Hypocoercivity and diffusion limit of a finite volume scheme for linear kinetic equations. *Math. Comp.* 89, no. 323, 1093–1133, 2020.
- [4] Krzysztof Bogdan, Andrzej St      , and Pawe   Sztonyk. Harnack inequality for stable processes on d-sets. *Studia Math.* 158, no. 2, 163–198, 2003.
- [5] Emeric Bouin, Jean Dolbeault, Laurent Lafleche, and Christian Schmeiser. Fractional Hypocoercivity. *arXiv preprint arXiv:1911.11020*, 2019.
- [6] Djalil Chafa    . Entropies, convexity, and functional inequalities: on Φ -entropies and Φ -Sobolev inequalities. *J. Math. Kyoto Univ.*, 44(2):325–363, 2004.
- [7] Nicolas Crouseilles, H       Hivert, and Mohammed Lemou. Numerical schemes for kinetic equations in the anomalous diffusion limit. Part I: The case of heavy-tailed equilibrium. *SIAM J. Sci. Comput.*, 38(2):A737–A764, 2016.
- [8] Guillaume Dujardin, Fr         H      , and Pauline Lafitte. Coercivity, hypocoercivity, exponential time decay and simulations for discrete Fokker-Planck equations. *Numer. Math.* 144, no. 3, 615–697, 2020.
- [9] Ivan Gentil and Cyril Imbert. The L      -Fokker-Planck equation: Φ -entropies and convergence to equilibrium. *Asymptot. Anal.*, 59(3-4):125–138, 2008.
- [10] Fr         H      . Introduction to hypocoercive methods and applications for simple linear inhomogeneous kinetic models. In *Lectures on the analysis of nonlinear partial differential equations. Part 5*, volume 5 of *Morningside Lect. Math.*, pages 119–147. Int. Press, Somerville, MA, 2018.
- [11] Fr         H      , Daniela Tonon, and Isabelle Tristani. Short time diffusion properties of inhomogeneous kinetic equations with fractional collision kernel. *arXiv preprint arXiv:1709.09943*, 2018.
- [12] Mateusz Kwa        . Ten equivalent definitions of the fractional Laplace operator. *Fract. Calc. Appl. Anal.*, 20(1):7–51, 2017.
- [13] Isabelle Tristani. Fractional Fokker-Planck equation. *Commun. Math. Sci.*, no. 5, 1243–1260, 2015.
- [14] C       Villani. Hypocoercivity. *Mem. Amer. Math. Soc.*, 202(950):iv+141, 2009.
- [15] Jian Wang. A simple approach to functional inequalities for non-local Dirichlet forms. *ESAIM, Probab. Stat.*, 18, 503–513, 2014.